

A CONSTRUCTIVE DEMONSTRATION OF THE UNIQUENESS OF THE CHRISTOFFEL SYMBOL

Victor Tapia¹

*Departamento de Física
Universidad Nacional de Colombia
Bogotá, Colombia*

The Christoffel symbol

$$\{\overset{\lambda}{\underset{\mu\nu}{\Gamma}}\}(\mathbf{g}) = \frac{1}{2} g^{\lambda\rho} (\partial_\mu g_{\nu\rho} + \partial_\nu g_{\mu\rho} - \partial_\rho g_{\mu\nu}), \quad (1)$$

is usually introduced (Spivak, 1975) as the connection which solves the metricity condition

$$\nabla_\lambda g_{\mu\nu} = 0. \quad (2)$$

On the other hand, whether the Christoffel symbol is the only connection which can be constructed from a symmetric second-rank tensor $g_{\mu\nu}$ remains an open question. In this note we exhibit a constructive demonstration of the uniqueness of the Christoffel symbol.

Let us start by reviewing some simple results of tensor calculus. Let \mathcal{M} be an n -dimensional differentiable manifold. Several geometric objects can be introduced over conveniently defined fibered bundles based on \mathcal{M} . In order to classify them we adopt a taxonomic approach, *cf.* (Visconti, 1992): a *tensor* is an object which transforms like a *tensor*, etc. The previous definition makes reference to the way in which an object transforms under changes of local coordinates. Let x^μ , $\mu = 1, 2, \dots, n$, and y^α , $\alpha = 1, 2, \dots, n$, be local coordinates on \mathcal{M} . Both sets are related by $y^\alpha = y^\alpha(x^\mu)$, and the differential form of this relation is

$$dy^\alpha = \left(\frac{\partial y^\alpha}{\partial x^\mu} \right) dx^\mu. \quad (3)$$

Due to the intrinsic function theorem, this relation tells us that y^α are functions of x^μ , $y^\alpha = y^\alpha(x^\mu)$. In order to express x^μ as functions of y^α , $x^\mu = x^\mu(y^\alpha)$, we need to invert eq. (3). This can be achieved if

$$\det \left(\frac{\partial y^\alpha}{\partial x^\mu} \right) \neq 0. \quad (4)$$

¹TAPIENS@CIENCIAS.CIENCIAS.UNAL.EDU.CO

If this is the case we have an inverse matrix $(\partial x/\partial y)$ and the relation (3) can be inverted to

$$dx^\mu = \left(\frac{\partial x^\mu}{\partial y^\alpha} \right) dy^\alpha. \quad (5)$$

Then $x^\mu = x^\mu(y^\alpha)$.

Now we are ready to proceed to our taxonomic classification of geometrical objects. Relation (3) provides the first example. Let us consider a set of functions $v^\mu(x)$ and $v^\alpha(y)$ which are related in the way given by (3)

$$v^\alpha(y) = \left(\frac{\partial y^\alpha}{\partial x^\mu} \right) v^\mu(x). \quad (6)$$

This defines a contravariant vector. Accordingly, a set of functions $v_\mu(x)$ and $v_\alpha(y)$ which are related by

$$v_\alpha(y) = \left(\frac{\partial x^\mu}{\partial y^\alpha} \right) v_\mu(x), \quad (7)$$

defines a covariant vector. Let us now consider the functions $\phi(x) = v_\mu(x)v^\mu(x)$ and $\phi(y) = v_\alpha(y)v^\alpha(y)$. From the relations above we easily obtain

$$\phi(y) = \phi(x). \quad (8)$$

This relation defines a scalar. The relations above can be extended to tensors of different rank and covariance. For example, a covariant second-rank tensor $g_{\mu\nu}(x)$ is an object which transforms as

$$g_{\alpha\beta}(y) = \left(\frac{\partial x^\mu}{\partial y^\alpha} \right) \left(\frac{\partial x^\nu}{\partial y^\beta} \right) g_{\mu\nu}(x). \quad (9)$$

Let us now consider the derivative of the scalar function (8). We obtain

$$\frac{\partial \phi}{\partial y^\alpha}(y) = \left(\frac{\partial x^\mu}{\partial y^\alpha} \right) \frac{\partial \phi}{\partial x^\mu}(x). \quad (10)$$

Therefore, the ordinary derivative of a scalar function is a vector. Next, let us consider the ordinary derivative of a covariant vector. From (7) we obtain

$$\frac{\partial v_\beta}{\partial y^\alpha}(y) = \left(\frac{\partial x^\mu}{\partial y^\alpha} \right) \left(\frac{\partial x^\nu}{\partial y^\beta} \right) \frac{\partial v_\nu}{\partial x^\mu}(x) + \left(\frac{\partial^2 x^\mu}{\partial y^\alpha \partial y^\beta} \right) v_\mu(x). \quad (11)$$

Therefore, this quantity is not a vector. Let us observe that covariance is broken by a term linear in v_μ . Let us therefore introduce a new derivative

$$\nabla_\mu v_\nu = \partial_\mu v_\nu - \Gamma_{\mu\nu}^\lambda v_\lambda, \quad (12)$$

where a term linear in v_μ is introduced to compensate the wrong behaviour of the last term in (11). Let us now impose that this quantity be a tensor. The result is that $\mathbf{\Gamma}$ must transform according to

$$\Gamma^\gamma_{\alpha\beta}(y) = \left(\frac{\partial x^\mu}{\partial y^\alpha}\right) \left(\frac{\partial x^\nu}{\partial y^\beta}\right) \left[\left(\frac{\partial y^\gamma}{\partial x^\lambda}\right) \Gamma^\lambda_{\mu\nu}(x) - \left(\frac{\partial^2 y^\gamma}{\partial x^\mu \partial x^\nu}\right) \right]. \quad (13)$$

It is clear that $\mathbf{\Gamma}$ is not a tensor; this is not unexpected since in order to compensate the non-tensor character of ∂v we need a non tensor object. $\nabla_\mu v_\nu$ is the covariant derivative, and $\mathbf{\Gamma}$ is the connection.

In Riemannian geometry the natural object is the metric tensor $g_{\mu\nu}$. The metric is not related to the connection. However, we may relate them through a metricity condition. The simplest metricity condition is

$$\nabla_\lambda^\Gamma g_{\mu\nu} = 0. \quad (14)$$

The relation (14) has the unique solution

$$\Gamma^\lambda_{\mu\nu} = \{\lambda_{\mu\nu}\}(\mathbf{g}) = \frac{1}{2} g^{\lambda\rho} (\partial_\mu g_{\nu\rho} + \partial_\nu g_{\mu\rho} - \partial_\rho g_{\mu\nu}), \quad (15)$$

which is known as the Christoffel symbol.

The Christoffel symbol is the connection which solves the metricity condition (14). It happens to be constructed from a second-rank tensor. However, there is a further question we may ask: which is the more general connection we can construct in terms of a second-rank tensor? We must therefore look for a more general scheme providing a most complete answer.

Let us therefore consider the problem of how to construct a connection $\Gamma^\lambda_{\mu\nu}$ only from a second-rank tensor $g_{\mu\nu}$. For this purpose let us remind that the connection is related to a derivation and therefore has dimensions corresponding to an inverse length, *i.e.*

$$\dim[\mathbf{\Gamma}] = L^{-1}. \quad (16)$$

On the other hand, the derivation is a linear operation. Therefore, the needed connection must be a linear combination of derivatives of $g_{\mu\nu}$, namely,

$$\Gamma^\lambda_{\mu\nu}(\mathbf{g}) = G^\lambda_{\mu\nu}{}^{\sigma\tau\rho}(\mathbf{g}, \mathbf{g}^{-1}) \partial_\sigma g_{\tau\rho}, \quad (17)$$

where \mathbf{G} is a function depending only on $g_{\mu\nu}$ and its inverse $g^{\mu\nu}$, and therefore it is a tensor. The transformation rule for $\mathbf{\Gamma}$ in (17) is

$$\begin{aligned} \Gamma^\gamma_{\alpha\beta}(y) &= G^\gamma_{\alpha\beta}{}^{\delta\epsilon\phi}(\mathbf{g}, \mathbf{g}^{-1}) \partial_\delta g_{\epsilon\phi} \\ &= G^\gamma_{\alpha\beta}{}^{\delta\epsilon\phi}(\mathbf{g}, \mathbf{g}^{-1}) \partial_\delta \left[\left(\frac{\partial x^\mu}{\partial y^\epsilon}\right) \left(\frac{\partial x^\nu}{\partial y^\phi}\right) g_{\mu\nu} \right] \end{aligned}$$

$$\begin{aligned}
&= \left(\frac{\partial y^\gamma}{\partial x^\lambda} \right) \left(\frac{\partial y^\alpha}{\partial x^\mu} \right) \left(\frac{\partial y^\beta}{\partial x^\nu} \right) \Gamma^\lambda{}_{\mu\nu}(x) \\
&\quad + 2 G_{\alpha\beta}^{\gamma\delta\epsilon\phi}(\mathbf{g}, \mathbf{g}^{-1}) \left(\frac{\partial^2 x^\mu}{\partial y^\delta \partial y^\epsilon} \right) \left(\frac{\partial x^\nu}{\partial y^\phi} \right) g_{\mu\nu} \\
&= \left(\frac{\partial y^\gamma}{\partial x^\lambda} \right) \left(\frac{\partial y^\alpha}{\partial x^\mu} \right) \left(\frac{\partial y^\beta}{\partial x^\nu} \right) \Gamma^\lambda{}_{\mu\nu}(x) \\
&\quad + 2 G_{\alpha\beta}^{\gamma\delta\epsilon\phi}(\mathbf{g}, \mathbf{g}^{-1}) g_{\phi\eta} \left(\frac{\partial y^\eta}{\partial x^\lambda} \right) \left(\frac{\partial^2 x^\lambda}{\partial y^\delta \partial y^\epsilon} \right). \tag{18}
\end{aligned}$$

Now we require that this object transform like a connection and if we compare with (13) we obtain

$$2 G_{\alpha\beta}^{\gamma\delta\epsilon\phi}(\mathbf{g}, \mathbf{g}^{-1}) g_{\phi\eta} \left(\frac{\partial y^\eta}{\partial x^\lambda} \right) \left(\frac{\partial^2 x^\lambda}{\partial y^\delta \partial y^\epsilon} \right) = \left(\frac{\partial y^\gamma}{\partial x^\lambda} \right) \left(\frac{\partial^2 x^\lambda}{\partial y^\alpha \partial y^\beta} \right). \tag{19}$$

Therefore

$$\left[4 G_{\alpha\beta}^{\gamma\delta\epsilon\phi}(\mathbf{g}, \mathbf{g}^{-1}) g_{\phi\eta} - \delta_\eta^\gamma (\delta_\alpha^\delta \delta_\beta^\epsilon + \delta_\beta^\delta \delta_\alpha^\epsilon) \right] \left(\frac{\partial y^\eta}{\partial x^\lambda} \right) \left(\frac{\partial^2 x^\lambda}{\partial y^\delta \partial y^\epsilon} \right) = 0. \tag{20}$$

What must be zero is the symmetric part of the square bracket and then we obtain

$$2 G_{\alpha\beta}^{\gamma\delta\epsilon\phi}(\mathbf{g}, \mathbf{g}^{-1}) g_{\phi\eta} + 2 G_{\alpha\beta}^{\gamma\epsilon\delta\phi}(\mathbf{g}, \mathbf{g}^{-1}) g_{\phi\eta} - \delta_\eta^\gamma (\delta_\alpha^\delta \delta_\beta^\epsilon + \delta_\beta^\delta \delta_\alpha^\epsilon) = 0. \tag{21}$$

Considering cyclic permutations of the indices $\delta\epsilon\phi$ we arrive to a set of equations which allow to determine \mathbf{G} . The solution is

$$\begin{aligned}
&G_{\alpha\beta}^{\gamma\delta\epsilon\phi}(\mathbf{g}, \mathbf{g}^{-1}) \\
&= \frac{1}{4} \left[g^{\gamma\phi} (\delta_\alpha^\delta \delta_\beta^\epsilon + \delta_\beta^\delta \delta_\alpha^\epsilon) + g^{\gamma\delta} (\delta_\alpha^\epsilon \delta_\beta^\phi + \delta_\beta^\epsilon \delta_\alpha^\phi) - g^{\gamma\epsilon} (\delta_\alpha^\delta \delta_\beta^\phi + \delta_\beta^\delta \delta_\alpha^\phi) \right]. \tag{22}
\end{aligned}$$

If we now replace this result in (16) we obtain that the connection is precisely the Christoffel symbol, namely eq. (15).

One can immediately check that

$$\nabla_\lambda^g g_{\mu\nu} \equiv 0, \tag{23}$$

where ∇^g is the covariant derivative constructed with the Christoffel symbol.

We have shown that the Christoffel symbol is the only connection which can be constructed from a symmetric second-rank tensor. The metricity condition (23) appears just as a side result and not as the starting point for the construction of the Christoffel symbol.

References

1. M. Spivak, *A comprehensive introduction to differential geometry* (Publish or Perish, Boston, 1975).
2. A. Visconti, *Introductory differential geometry for physicists* (World Scientific, River Edge, 1992).